
Product Name: DGK- η Rabbit Polyclonal Antibody**Catalog #: APRab09951**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	IHC,ICC/IF,ELISA
Reactivity	Human,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000

Molecular Weight

Antigen Information

Gene Name	DGKH
Alternative Names	DGKH; Diacylglycerol kinase eta; DAG kinase eta; Diglyceride kinase eta; DGK-eta
Gene ID	160851.0
SwissProt ID	Q86XP1
Immunogen	The antiserum was produced against synthesized peptide derived from human DGKH. AA range:771-820

Background


diacylglycerol kinase eta(DGKH) Homo sapiens This gene encodes a member of the diacylglycerol kinase (DGK) enzyme

family. Members of this family are involved in regulating intracellular concentrations of diacylglycerol and phosphatidic acid. Variation in this gene has been associated with bipolar disorder. Alternatively spliced transcript variants have been identified. [provided by RefSeq, Jul 2014],catalytic activity:ATP + 1,2-diacylglycerol = ADP + 1,2-diacyl-sn-glycerol 3-phosphate.,function:Phosphorylates diacylglycerol (DAG) to generate phosphatidic acid (PA).,PTM:Phosphorylated; does not inhibit catalytic activity.,similarity:Belongs to the eukaryotic diacylglycerol kinase family.,similarity:Contains 1 DAGKc domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 SAM (sterile alpha motif) domain.,similarity:Contains 2 phorbol-ester/DAG-type zinc fingers.,subcellular location:Translocated from the cytoplasm to endosomes in response to stress stimuli. Isoform 2 is rapidly relocated back to the cytoplasm upon removal of stress stimuli, whereas isoform 1 exhibits sustained endosomal association.,subunit:Isoform 1 forms homooligomers through the SAM domain. Isoform 1 is also able to form heterooligomers with SAM domain-containing isoforms of DGKD. Oligomerization of isoform 1 inhibits its catalytic activity.,tissue specificity:Isoform 2 is ubiquitously expressed. Isoform 1 is expressed only in testis, kidney and colon.,


Research Area

Glycerolipid metabolism;Glycerophospholipid metabolism;Phosphatidylinositol signaling system;

Image Data



Immunofluorescence analysis of HeLa cells, using DGKH Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using DGKH Antibody. The picture on the right is blocked with the synthesized peptide.