
Product Name: FA2H Rabbit Polyclonal Antibody**Catalog #: APRab10744**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ELISA 1:10000-1:20000
Molecular Weight	55kDa

Antigen Information

Gene Name	FA2H
Alternative Names	Fatty acid 2-hydroxylase (EC 1.-.-.) (Fatty acid alpha-hydroxylase)
Gene ID	79152.0
SwissProt ID	Q7L5A8
Immunogen	Synthesized peptide derived from human FA2H. at AA range: 101-150


Background

This gene encodes a protein that catalyzes the synthesis of 2-hydroxysphingolipids, a subset of sphingolipids that contain 2-hydroxy fatty acids. Sphingolipids play roles in many cellular processes and their structural diversity arises from modification of

the hydrophobic ceramide moiety, such as by 2-hydroxylation of the N-acyl chain, and the existence of many different head groups. Mutations in this gene have been associated with leukodystrophy dysmyelinating with spastic paraparesis with or without dystonia.[provided by RefSeq, Mar 2010],cofactor:Iron.,disease:Defects in FA2H are the cause of leukodystrophy dysmyelinating with spastic paraparesis with or without dystonia (DLDSP) [MIM:612443]. The disorder consists of a progressive neurologic disease manifested by spasticity, disordered tonic of muscle, and white matter degeneration.,domain:The histidine box domains may contain the active site and/or be involved in metal ion binding.,function:Required for alpha-hydroxylation of free fatty acids and the formation of alpha-hydroxylated sphingolipids.,induction:Up-regulated during keratinocyte differentiation.,similarity:Belongs to the SCS7 family.,similarity:Contains 1 cytochrome b5 heme-binding domain.,tissue specificity:Detected in differentiating cultured keratinocytes (at protein level). Detected in epidermis and cultured keratinocytes. Highly expressed in brain and colon. Detected at lower levels in testis, prostate, pancreas and kidney.,

Research Area

Image Data



Western Blot analysis of mouse-heart cells using FA2H Rabbit Polyclonal Antibody diluted at 1:2000 (4°C overnight) . Secondary antibody : Goat Anti-rabbit IgG IRDye 800 (diluted at 1:5000, 25°C, 1 hour)