
Product Name: FBP2 Rabbit Polyclonal Antibody**Catalog #: APRab10860**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ELISA 1:5000-1:20000
Molecular Weight	73kDa

Antigen Information

Gene Name	KHSRP
Alternative Names	KHSRP; FUBP2; Far upstream element-binding protein 2; FUSE-binding protein 2; KH type-splicing regulatory protein; KSRP; p75
Gene ID	8570.0
SwissProt ID	Q92945
Immunogen	Synthesized peptide derived from the N-terminal region of human FBP2.


Background

The KHSRP gene encodes a multifunctional RNA-binding protein implicated in a variety of cellular processes, including


transcription, alternative pre-mRNA splicing, and mRNA localization (Min et al., 1997 [PubMed 9136930]; Gherzi et al., 2004 [PubMed 15175153]).[supplied by OMIM, Apr 2010],function:Binds to the dendritic targeting element and may play a role in mRNA trafficking (By similarity). Part of a ternary complex that binds to the downstream control sequence (DCS) of the pre-mRNA. Mediates exon inclusion in transcripts that are subject to tissue-specific alternative splicing. May interact with single-stranded DNA from the far-upstream element (FUSE). May activate gene expression. Also involved in degradation of inherently unstable mRNAs that contain AU-rich elements (AREs) in their 3'-UTR, possibly by recruiting degradation machinery to ARE-containing mRNAs.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the KHSRP family.,similarity:Contains 4 KH domains.,subcellular location:A small proportion is also found in the cytoplasm of neuronal cell bodies and dendrites.,subunit:Part of a ternary complex containing FUBP2, PTBP1, PTBP2 and HNRPH1. Interacts with PARN. Also interacts with APC which is part of the apolipoprotein B mRNA editing enzyme complex. The interaction is not required for activity of the catalytic component, APOBEC1, but may confer stability on the complex.,tissue specificity:Detected in neural and non-neural cell lines.,

Research Area

Image Data



Western Blot analysis of various cells using FBP2 Polyclonal Antibody



Western Blot analysis of HuvEc cells using FBP2 Polyclonal Antibody