

Product Name: FKBP1B Rabbit Polyclonal Antibody**Catalog #: APRab11003**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	IHC,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio IHC 1:50-1:200,ELISA 1:5000-1:20000

Molecular Weight

Antigen Information


Gene Name	FKBP1B
Alternative Names	Peptidyl-prolyl cis-trans isomerase FKBP1B (PPIase FKBP1B;EC 5.2.1.8;12.6 kDa FK506-binding protein;12.6 kDa FKBP;FKBP-12.6;FK506-binding protein 1B;FKBP-1B;Immunophilin FKBP12.6;Rotamase;h-FKBP-12)
Gene ID	2281.0
SwissProt ID	P68106
Immunogen	Synthesized peptide derived from human FKBP1B AA range: 35-85

Background

The protein encoded by this gene is a member of the immunophilin protein family, which play a role in immunoregulation and basic cellular processes involving protein folding and trafficking. This encoded protein is a cis-trans prolyl isomerase that binds the immunosuppressants FK506 and rapamycin. It is highly similar to the FK506-binding protein 1A. Its physiological role is thought to be in excitation-contraction coupling in cardiac muscle. There are two alternatively spliced transcript variants of this gene encoding different isoforms. [provided by RefSeq, Jul 2008],catalytic activity:Peptidylproline (omega=180) = peptidylproline (omega=0),enzyme regulation:Inhibited by both FK506 and rapamycin.,function:Associates with the ryanodine receptor (RyR-2) in cardiac muscle sarcoplasmic reticulum and may play a unique physiological role in excitation-contraction coupling in cardiac muscle. There are four molecules of FKBP12.6 per heart muscle RyR. Has the potential to contribute to the immunosuppressive and toxic effects of FK506 and rapamycin. PPIases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides.,similarity:Belongs to the FKBP-type PPIase family. FKBP1 subfamily.,similarity:Contains 1 PPIase FKBP-type domain.,tissue specificity:Ubiquitous for both isoforms with highest levels in brain and thymus.,

Research Area

Image Data



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200 (4° overnight) . 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .