
Product Name: FBNP4 Rabbit Polyclonal Antibody**Catalog #: APRab11053**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ELISA 1:5000-1:20000
Molecular Weight	111kDa

Antigen Information

Gene Name	FBNP4 FBP30 KIAA1014
Alternative Names	
Gene ID	23360.0
SwissProt ID	Q8N3X1
Immunogen	Synthesized peptide derived from part region of human protein


Background

domain:These WW domains interact with Arg/Gly-rich-flanked Pro-rich domains found in several WW domain-binding proteins (WBPs). The N-terminal WW domain has the greater ligand-binding ability.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 2 WW domains.,subunit:Binds FMN1. Interacts with the Arg/Gly-rich-flanked Pro-

rich of KHDRBS1/SAM68. Arginine methylation in these regions has no effect on this binding.,domain:These WW domains interact with Arg/Gly-rich-flanked Pro-rich domains found in several WW domain-binding proteins (WBPs). The N-terminal WW domain has the greater ligand-binding ability.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 2 WW domains.,subunit:Binds FMN1. Interacts with the Arg/Gly-rich-flanked Pro-rich of KHDRBS1/SAM68. Arginine methylation in these regions has no effect on this binding.,

Research Area

Image Data



Western Blot analysis of HeLa lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000