# **Product Name: Glut5 Rabbit Polyclonal Antibody**

**C** EnkiLife

Catalog #: APRab11505

## **Summary**

**Production Name** Glut5 Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

Host Rabbit
Application WB,ELISA

**Reactivity** Human,Rat,Mouse

### **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer** 

preservative N.

**Purification** Affinity purification

### **Immunogen**

Gene Name SLC2A5

SLC2A5; GLUT5; Solute carrier family 2, facilitated glucose transporter member 5; Alternative Names

Fructose transporter; Glucose transporter type 5, small intestine; GLUT-5

**Gene ID** 6518.0

P22732.The antiserum was produced against synthesized peptide derived from the N-SwissProt ID

terminal region of human SLC2A5. AA range:31-80

## **Application**

**Dilution Ratio** WB 1:500-1:2000, ELISA 1:20000.Not yet tested in other applications.

Molecular Weight 55kDa

# **Product Name: Glut5 Rabbit Polyclonal Antibody**

Catalog #: APRab11505




# **Background**


The protein encoded by this gene is a fructose transporter responsible for fructose uptake by the small intestine. The encoded protein also is necessary for the increase in blood pressure due to high dietary fructose consumption. [provided by RefSeq, Jun 2016], function: Cytochalasin B-sensitive carrier. Seems to function primarily as a fructose transporter., induction: By forskolin (in Caco-2 cells)., mass spectrometry: PubMed: 11840567, similarity: Belongs to the major facilitator superfamily. Sugar transporter (TC 2.A.1.1) family. Glucose transporter subfamily., tissue specificity: Expressed in small intestine, and at much lower levels in kidney, skeletal muscle, and adipose tissue.,

#### **Research Area**

### **Image Data**



Western blot analysis of lysate from K562 cells, using SLC2A5 Antibody.



Western Blot analysis of K562 cells using Glut5 Polyclonal Antibody. Secondary antibody was diluted at 1:20000

#### Note

For research use only.