
Product Name: IL-1F7 Rabbit Polyclonal Antibody**Catalog #: APRab12521**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	IHC,ICC/IF,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio IHC 1:50-1:200,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000

Molecular Weight

Antigen Information


Gene Name	IL37 Interleukin-37 (FIL1 zeta;IL-1X;Interleukin-1 family member 7;IL-1F7;Interleukin-1 homolog
Alternative Names	4;IL-1H;IL-1H4;Interleukin-1 zeta;IL-1 zeta;Interleukin-1-related protein;IL-1RP1;Interleukin-23;IL-37)
Gene ID	27178.0
SwissProt ID	Q9NZH6
Immunogen	Synthetic peptide from human protein at AA range: 101-150

Background


The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine can bind to, and may be a ligand for interleukin 18 receptor (IL18R1/IL-1R α). This cytokine also binds to interleukin 18 binding protein (IL18BP), an inhibitory binding protein of interleukin 18 (IL18), and subsequently forms a complex with IL18 receptor beta subunit, and through which it inhibits the activity of IL18. This gene along with eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. Five alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008],function: Binds to interleukin-18 receptor (IL-18R) receptor but not to IL-1 receptor. Could be a new player in the inflammatory and immune responses mediated by the IL-18/IL-18R axis.,induction: By phorbol ester (PMA) in different cell lines.,miscellaneous: The sequence shown here is derived from an EMBL/GenBank/DDBJ third party annotation (TPA) entry.,online information: Interleukin-1 entry,similarity: Belongs to the IL-1 family.,tissue specificity: Isoform A, isoform B and isoform C are expressed in testis, colon, placenta, lung and lymph node. Isoform D and isoform E were found only in testis and bone marrow. Whereas only isoform A is found in brain, only isoform B in kidney and only isoform C in heart.,

Research Area


Image Data




Immunohistochemical analysis of paraffin-embedded human-small-intestine, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-small-intestine, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-colon-cancer, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-colon-cancer, antibody was diluted at 1:200