

Product Name: IL-28R α Rabbit Polyclonal Antibody**Catalog #: APRab12545**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ELISA 1:20000-1:40000
Molecular Weight	57kDa

Antigen Information

Gene Name	IL28RA
Alternative Names	IL28RA; IFNLR1; LICR2; Interleukin-28 receptor subunit alpha; IL-28 receptor subunit alpha; IL-28R-alpha; IL-28RA; Cytokine receptor class-II member 12; Cytokine receptor family 2 member 12;CRF2-12; Interferon lambda receptor 1; IFN-lambda receptor 1; IFN-lambda-R1; Likely interleukin or cytokine receptor 2; LICR2
Gene ID	163702.0
SwissProt ID	Q8IU57
Immunogen	Synthesized peptide derived from the Internal region of human IL-28R α .


Background

The protein encoded by this gene belongs to the class II cytokine receptor family. This protein forms a receptor complex with interleukine 10 receptor, beta (IL10RB). The receptor complex has been shown to interact with three closely related cytokines, including interleukin 28A (IL28A), interleukin 28B (IL28B), and interleukin 29 (IL29). The expression of all three cytokines can be induced by viral infection. The cells overexpressing this protein have been found to have enhanced responses to IL28A and IL29, but decreased response to IL28B. Three alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008],function:The IL28RA/IL10RB dimer is a receptor for IL28A, IL28B and IL29. The ligand/receptor complex seems to signal through the Jak-STAT pathway.,similarity:Belongs to the type II cytokine receptor family.,similarity:Contains 1 fibronectin type-III domain.,subunit:Heterodimer with IL10RB.,tissue specificity:Widely expressed.,

Research Area

Cytokine-cytokine receptor interaction;Jak_STAT;

Image Data



Western Blot analysis of extracts from K562 cells, using IL-28R α Polyclonal Antibody.. Secondary antibody was diluted at 1:20000