
Product Name: IL-31R α Rabbit Polyclonal Antibody**Catalog #: APRab12551**

For research use only.

Summary

| | |
|----------------------|---|
| Description | Rabbit polyclonal Antibody |
| Host | Rabbit |
| Application | WB,ELISA |
| Reactivity | Human,Rat,Mouse |
| Conjugation | Unconjugated |
| Modification | Unmodified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Form | Liquid |
| Concentration | 1mg/ml |
| Storage | Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles. |
| Shipping | Ice bags |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| Purification | Affinity purification |

Application

| | |
|-------------------------|---------------------------------------|
| Dilution Ratio | WB 1:500-1:2000,ELISA 1:20000-1:40000 |
| Molecular Weight | 82kDa |

Antigen Information


| | |
|--------------------------|---|
| Gene Name | IL31RA |
| Alternative Names | IL31RA; CRL3; GPL; Interleukin-31 receptor subunit alpha; IL-31 receptor subunit alpha; IL-31R subunit alpha; IL-31R-alpha; IL-31RA; Cytokine receptor-like 3; GLM-R; hGLM-R; Gp130-like monocyte receptor; Gp130-like receptor; ZcytoR17 |
| Gene ID | 133396.0 |
| SwissProt ID | Q8NI17 |
| Immunogen | Synthesized peptide derived from the Internal region of human IL-31R α . |

Background

The protein encoded by this gene belongs to the type I cytokine receptor family. This receptor, with homology to gp130, is expressed on monocytes, and is involved in IL-31 signaling via activation of STAT-3 and STAT-5. It functions either as a monomer, or as part of a receptor complex with oncostatin M receptor (OSMR). Several alternatively spliced transcript variants encoding different isoforms have been noted for this gene.[provided by RefSeq, Jun 2011],function:Associates with OSMR to form the interleukin-31 receptor which activates STAT3 and to a lower extent STAT1 and STAT5. May function in skin immunity.,induction:Up-regulated in lesional keratinocytes of patients with atopic dermatitis. Up-regulated by IFNG and lipopolysaccharides.,PTM:N-glycosylated.,similarity:Belongs to the type I cytokine receptor family. Type 2 subfamily.,similarity:Contains 5 fibronectin type-III domains.,subunit:Heterodimer with OSMR. Interacts with JAK1 and STAT3.,tissue specificity:Expressed at low levels in testis, ovary, brain, prostate, placenta, thymus, bone marrow, trachea and skin. Detected in all of the myelomonocytic lineage. Expressed in CD14- and CD56-positive blood cells and by macrophages (at protein level),,

Research Area

Image Data



Western Blot analysis of extracts from Jurkat cells, using IL-31R Polyclonal Antibody. Secondary antibody was diluted at 1:20000