

Product Name: IL-5 Rabbit Polyclonal Antibody**Catalog #: APRab12562**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:10000
Molecular Weight	30kDa

Antigen Information

Gene Name	IL5
Alternative Names	IL5; Interleukin-5; IL-5; B-cell differentiation factor I; Eosinophil differentiation factor; T-cell replacing factor; TRF
Gene ID	3567.0
SwissProt ID	P05113
Immunogen	The antiserum was produced against synthesized peptide derived from human IL-5. AA range:43-92


Background

This gene encodes a cytokine that acts as a growth and differentiation factor for both B cells and eosinophils. The encoded cytokine plays a major role in the regulation of eosinophil formation, maturation, recruitment and survival. The increased production of this cytokine may be related to pathogenesis of eosinophil-dependent inflammatory diseases. This cytokine functions by binding to its receptor, which is a heterodimer, whose beta subunit is shared with the receptors for interleukine 3 (IL3) and colony stimulating factor 2 (CSF2/GM-CSF). This gene is located on chromosome 5 within a cytokine gene cluster which includes interleukin 4 (IL4), interleukin 13 (IL13), and CSF2 . This gene, IL4, and IL13 may be regulated coordinately by long-range regulatory elements spread over 120 kilobases on chromosome 5q31. [provided by RefSeq, Jul 2013],function:Factor that induces terminal differentiation of late-developing B-cells to immunoglobulin secreting cells.,online information:Interleukin-5 entry,similarity:Belongs to the IL-5 family.,subunit:Homodimer; disulfide-linked.,


Research Area

Cytokine-cytokine receptor interaction;Jak_STAT;Hematopoietic cell lineage;T_Cell_Receptor;Fc epsilon RI;Intestinal immune network for IgA production;Asthma;Autoimmune thyroid disease;Allograft rejection;


Image Data



Immunohistochemistry analysis of IL-5 antibody in paraffin-embedded human breast carcinoma tissue.



Western blot analysis of lysate from HT-29 cells, using IL-5 antibody.



Western Blot analysis of various cells using IL-5 Polyclonal Antibody