

Product Name: IL-9R Rabbit Polyclonal Antibody**Catalog #: APRab12575**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ELISA 1:5000-1:20000
Molecular Weight	58kDa

Antigen Information

Gene Name	IL9R
Alternative Names	IL9R; Interleukin-9 receptor; IL-9 receptor; IL-9R; CD129
Gene ID	3581.0
SwissProt ID	Q01113
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human IL9R. AA range:201-250

Background


The protein encoded by this gene is a cytokine receptor that specifically mediates the biological effects of interleukin 9 (IL9).

The functional IL9 receptor complex requires this protein as well as the interleukin 2 receptor, gamma (IL2RG), a common gamma subunit shared by the receptors of many different cytokines. The ligand binding of this receptor leads to the activation of various JAK kinases and STAT proteins, which connect to different biologic responses. This gene is located at the pseudoautosomal regions of X and Y chromosomes. Genetic studies suggested an association of this gene with the development of asthma. Multiple pseudogenes on chromosome 9, 10, 16, and 18 have been described. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2008],domain:The box 1 motif is required for JAK interaction and/or activation.,domain:The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding.,function:This is a receptor for interleukin-9.,miscellaneous:The gene encoding for this protein is located in the pseudoautosomal region 2 (PAR2) of X and Y chromosomes.,similarity:Belongs to the type I cytokine receptor family. Type 4 subfamily.,similarity:Contains 1 fibronectin type-III domain.,


Research Area

Cytokine-cytokine receptor interaction;Jak_STAT;Hematopoietic cell lineage;

Image Data



Western Blot analysis of KB, NIH-3T3 cells using IL-9R Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Western Blot analysis of KB cells using IL-9R Polyclonal Antibody. Secondary antibody was diluted at 1:20000