

Summary

Production Name	LIMK-2 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	LIMK2
Alternative Names	LIMK2; LIM domain kinase 2; LIMK-2
Gene ID	3985.0
SwissProt ID	P53671.The antiserum was produced against synthesized peptide derived from human LIMK2. AA range:249-298

Application

Dilution Ratio	WB 1:500-1:2000, IHC-P 1:100-1:300, IF-P/IF-F/ICC/IF 1:200-1:1000, ELISA 1:40000.Not yet tested in other applications.
Molecular Weight	72kDa


Background

There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. The protein encoded by this gene is phosphorylated and activated by ROCK, a downstream effector of Rho, and the encoded protein, in turn, phosphorylates cofilin, inhibiting its actin-depolymerizing activity. It is thought that this pathway contributes to Rho-induced reorganization of the actin cytoskeleton. At least three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008], catalytic activity: ATP + a protein = ADP + a phosphoprotein., function: Displays serine/threonine-specific phosphorylation of myelin basic protein and histone (MBP) in vitro., PTM: Phosphorylated on serine and/or threonine residues by ROCK1., similarity: Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family., similarity: Contains 1 PDZ (DHR) domain., similarity: Contains 1 protein kinase domain., similarity: Contains 2 LIM zinc-binding domains., subcellular location: Isoform LIMK2a is distributed in the cytoplasm and the nucleus., subcellular location: Isoform LIMK2b occurs mainly in the cytoplasm and is scarcely translocated to the nucleus., subunit: Binds ROCK1 and LKAP. Interacts with PARD3. Interacts with NISCH., tissue specificity: Highest expression in the placenta; moderate level in liver, lung, kidney, and pancreas. LIMK2a is found to be more abundant than LIMK2b in liver, colon, stomach, and spleen, while in brain, kidney, and placenta LIMK2b is the dominant form. In adult lung, both LIMK2a and LIMK2b is nearly equally observed.,

Research Area


Axon guidance; Fc gamma R-mediated phagocytosis; Regulates Actin and Cytoskeleton;

Image Data




Western blot analysis of lysates from NIH/3T3 cells, treated with PMA 125ng/ml 30', using LIMK2 Antibody. The lane on the right is blocked with the synthesized peptide.

Product Name: LIMK-2 Rabbit Polyclonal Antibody
Catalog #: APRab13315



Western blot analysis of the lysates from RAW264.7 cells using LIMK2 antibody.



Western Blot analysis of 3T3 cells using LIMK-2 Polyclonal Antibody

Note

For research use only.