## **Product Name: LPHN3 Rabbit Polyclonal Antibody**

Catalog #: APRab13392



## **Summary**

Production Name LPHN3 Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

Host Rabbit
Application WB,ELISA

**Reactivity** Human, Mouse, Rat

## **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

Clonality Polyclonal Form Liquid

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

**Buffer** Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.

**Purification** Affinity purification

## **Immunogen**

Gene Name LPHN3 KIAA0768 LEC3

**Alternative Names** 

**Gene ID** 23284.0

**SwissProt ID** Q9HAR2. Synthesized peptide derived from part region of human protein

## **Application**

**Dilution Ratio** WB 1:500-2000, ELISA 1:5000-20000

Molecular Weight 159kDa

## **Background**

This gene encodes a member of the latrophilin subfamily of G-protein coupled receptors (GPCR). Latrophilins may function in both cell adhesion and signal transduction. In experiments with non-human species, endogenous proteolytic cleavage

# Product Name: LPHN3 Rabbit Polyclonal Antibody Catalog #: APRab13392



within a cysteine-rich GPS (G-protein-coupled-receptor proteolysis site) domain resulted in two subunits (a large extracellular N-terminal cell adhesion subunit and a subunit with substantial similarity to the secretin/calcitonin family of GPCRs) being non-covalently bound at the cell membrane. [provided by RefSeq, Jul 2008],PTM:Proteolytically cleaved into 2 subunits, an extracellular subunit and a seven-transmembrane subunit.,similarity:Belongs to the G-protein coupled receptor 2 family. LN-TM7 subfamily.,similarity:Contains 1 GPS domain.,similarity:Contains 1 olfactomedin-like domain.,similarity:Contains 1 SUEL-type lectin domain.,subunit:Forms a heterodimer, consisting of a large extracellular region (p120) non-covalently linked to a seven-transmembrane moiety (p85).,

### **Research Area**

### **Image Data**



Western Blot analysis of HEK293 lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:1000

#### Note

For research use only.