

Product Name: Mac-2BP Rabbit Polyclonal Antibody
Catalog #: APRab13541



Summary

Production Name	Mac-2BP Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	LGALS3BP M2BP
Alternative Names	Galectin-3-binding protein (Basement membrane autoantigen p105) (Lectin galactoside-binding soluble 3-binding protein) (Mac-2-binding protein) (MAC2BP) (Mac-2 BP) (Tumor-associated antigen 90K)
Gene ID	3959.0
SwissProt ID	Q08380. Synthetic peptide from human protein at AA range: 100-160

Application

Dilution Ratio	WB 1:500-2000, ELISA 1:10000-20000
Molecular Weight	65kDa

Product Name: Mac-2BP Rabbit Polyclonal Antibody
Catalog #: APRab13541




Background

The galectins are a family of beta-galactoside-binding proteins implicated in modulating cell-cell and cell-matrix interactions. LGALS3BP has been found elevated in the serum of patients with cancer and in those infected by the human immunodeficiency virus (HIV). It appears to be implicated in immune response associated with natural killer (NK) and lymphokine-activated killer (LAK) cell cytotoxicity. Using fluorescence in situ hybridization the full length 90K cDNA has been localized to chromosome 17q25. The native protein binds specifically to a human macrophage-associated lectin known as Mac-2 and also binds galectin 1. [provided by RefSeq, Jul 2008],function:Promotes intergrin-mediated cell adhesion. May stimulate host defense against viruses and tumor cells.,similarity:Contains 1 BACK (BTB/Kelch associated) domain.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 1 SRCR domain.,subunit:Homodimers and homomultimers. The multimers form ring-like structures with a diameter of 30-40 nm. Binds LGALS1 and LGALS3. Binds ITGB1, COL4A1, COL5A1, COL6A1, FN1 and NID.,tissue specificity:Ubiquitous. Detected in body fluids such as semen, milk, serum, tears, saliva and urine. Expressed by keratinocytes and fibroblasts.,

Research Area

Image Data



Western blot analysis of human blood lysate, antibody was diluted at 1000. Secondary antibody was diluted at 1:20000

Note

For research use only.