
Product Name: Manic Fringe Rabbit Polyclonal Antibody**Catalog #: APRab13619**

For research use only.

Summary

| | |
|----------------------|---|
| Description | Rabbit polyclonal Antibody |
| Host | Rabbit |
| Application | WB,ELISA |
| Reactivity | Human,Mouse,Rat |
| Conjugation | Unconjugated |
| Modification | Unmodified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Form | Liquid |
| Concentration | 1mg/ml |
| Storage | Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles. |
| Shipping | Ice bags |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| Purification | Affinity purification |

Application

| | |
|-------------------------|--------------------------------------|
| Dilution Ratio | WB 1:500-1:2000,ELISA 1:5000-1:10000 |
| Molecular Weight | 38kDa |

Antigen Information

| | |
|--------------------------|---|
| Gene Name | MFNG |
| Alternative Names | MFNG; Beta-1; 3-N-acetylglucosaminyltransferase manic fringe; O-fucosylpeptide 3-beta-N-acetylglucosaminyltransferase |
| Gene ID | 4242.0 |
| SwissProt ID | O00587 |
| Immunogen | The antiserum was produced against synthesized peptide derived from human MFNG. AA range:61-110 |


Background

This gene is a member of the fringe gene family which also includes radical and lunatic fringe genes. They all encode evolutionarily conserved secreted proteins that act in the Notch receptor pathway to demarcate boundaries during embryonic development. While their genomic structure is distinct from other glycosyltransferases, fringe proteins have a fucose-specific beta-1,3-N-acetylglucosaminyltransferase activity that leads to elongation of O-linked fucose residues on Notch, which alters Notch signaling. [provided by RefSeq, Oct 2009],catalytic activity:Transfers a beta-D-GlcNAc residue from UDP-D-GlcNAc to the fucose residue of a fucosylated protein acceptor.,function:Glycosyltransferase involved in the elongation of O-linked ligands to activate Notch signaling. Possesses fucose-specific beta-1,3-N-acetylglucosaminyltransferase activity.,online information:Beta-1,3-N-acetylglucosaminyltransferase manic fringe,online information:GlycoGene database,similarity:Belongs to the glycosyltransferase 31 family,.

Research Area

Notch;

Image Data



Western blot analysis of the lysates from HepG2 cells using MFNG antibody.