Product Name: MC2-R Rabbit Polyclonal Antibody

Catalog #: APRab13697



Summary

Production Name MC2-R Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit

Application WB,ELISA,IHC-P **Reactivity** Human,Mouse

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG

ClonalityPolyclonalFormLiquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer**

preservative N.

Purification Affinity purification

Immunogen

Gene Name MC2R

MC2R; ACTHR; Adrenocorticotropic hormone receptor; ACTH receptor; ACTH-R; Alternative Names

Adrenocorticotropin receptor; Melanocortin receptor 2; MC2-R

Gene ID 4158.0

Q01718.The antiserum was produced against synthesized peptide derived from human SwissProt ID

ACTHR. AA range:248-297

Application

Dilution Ratio WB 1:500-2000, IHC-P 1:50-300, ELISA 2000-20000

Molecular Weight 34kDa

Product Name: MC2-R Rabbit Polyclonal Antibody

Catalog #: APRab13697




Background

MC2R encodes one member of the five-member G-protein associated melanocortin receptor family. Melanocortins (melanocyte-stimulating hormones and adrenocorticotropic hormone) are peptides derived from pro-opiomelanocortin (POMC). MC2R is selectively activated by adrenocorticotropic hormone, whereas the other four melanocortin receptors recognize a variety of melanocortin ligands. Mutations in MC2R can result in familial glucocorticoid deficiency. Alternate transcript variants have been found for this gene. [provided by RefSeq, May 2014], disease:Defects in MC2R are the cause of glucocorticoid deficiency type 1 (GCCD1) [MIM:202200]; also known as familial glucocorticoid deficiency type 1 (FGD1). GCCD1 is an autosomal recessive disorder due to congenital insensitivity or resistance to adrenocorticotropin (ACTH). It is characterized by progressive primary adrenal insufficiency, without mineralocorticoid deficiency.,function:Receptor for ACTH. This receptor is mediated by G proteins (G(s)) which activate adenylate cyclase, similarity:Belongs to the G-protein coupled receptor 1 family, subunit:Interacts with FALP/MRAP, tissue specificity:Melanocytes and corticoadrenal tissue,


Research Area

Neuroactive ligand-receptor interaction;

Image Data




Immunofluorescence analysis of MCF7 cells, using ACTHR Antibody. The picture on the right is blocked with the synthesized




Product Name: MC2-R Rabbit Polyclonal Antibody Catalog #: APRab13697



Western blot analysis of lysates from HeLa cells, using ACTHR Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using ACTHR antibody.



Western Blot analysis of various cells using MC2-R Polyclonal Antibody

Note

For research use only.