
Product Name: ME2 Rabbit Polyclonal Antibody**Catalog #: APRab13763**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:20000-1:40000
Molecular Weight	65kDa

Antigen Information

Gene Name	ME2
Alternative Names	ME2; NAD-dependent malic enzyme; mitochondrial; NAD-ME; Malic enzyme 2
Gene ID	4200.0
SwissProt ID	P23368
Immunogen	The antiserum was produced against synthesized peptide derived from human ME2. AA range:201-250

Background


This gene encodes a mitochondrial NAD-dependent malic enzyme, a homotetrameric protein, that catalyzes the oxidative

decarboxylation of malate to pyruvate. It had previously been weakly linked to a syndrome known as Friedreich ataxia that has since been shown to be the result of mutation in a completely different gene. Certain single-nucleotide polymorphism haplotypes of this gene have been shown to increase the risk for idiopathic generalized epilepsy. Alternatively spliced transcript variants encoding different isoforms found for this gene. [provided by RefSeq, Dec 2009],catalytic activity:(S)-malate + NAD(+) = pyruvate + CO(2) + NADH.,cofactor:Divalent metal cations. Prefers magnesium or manganese.,enzyme regulation:Subject to allosteric activation by fumarate.,miscellaneous:This isoenzyme can also use NADP(+) but is more effective with NAD(+).,similarity:Belongs to the malic enzymes family.,subunit:Homotetramer.,


Research Area

Pyruvate metabolism;

Image Data



Immunohistochemistry analysis of paraffin-embedded human placenta tissue, using ME2 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from RAW264.7 and LOVO cells, using ME2 Antibody. The lane on the right is blocked with the synthesized peptide.