

**Product Name: MED11 Rabbit Polyclonal Antibody****Catalog #: APRab13769**

For research use only.

**Summary**

|                      |   |
|----------------------|---|
| <b>Description</b>   | Rabbit polyclonal Antibody  |
| <b>Host</b>          | Rabbit  |
| <b>Application</b>   | WB,ELISA  |
| <b>Reactivity</b>    | Human,Mouse   |
| <b>Conjugation</b>   | Unconjugated  |
| <b>Modification</b>  | Unmodified  |
| <b>Isotype</b>       | IgG   |
| <b>Clonality</b>     | Polyclonal  |
| <b>Form</b>          | Liquid  |
| <b>Concentration</b> | 1mg/ml  |
| <b>Storage</b>       | Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles. |
| <b>Shipping</b>      | Ice bags  |
| <b>Buffer</b>        | Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.   |
| <b>Purification</b>  | Affinity purification   |

**Application**

|                         |                                      |
|-------------------------|--------------------------------------|
| <b>Dilution Ratio</b>   | WB 1:500-1:2000,ELISA 1:5000-1:20000 |
| <b>Molecular Weight</b> | 12kDa                                |

**Antigen Information**

|                          |  |
|--------------------------|--|
| <b>Gene Name</b>         | MED11 HSPC296  |
| <b>Alternative Names</b> |  |
| <b>Gene ID</b>           | 400569.0   |
| <b>SwissProt ID</b>      | Q9P086   |
| <b>Immunogen</b>         | Synthesized peptide derived from human protein . at AA range: 30-110 |


**Background**

MED11 is a component of the Mediator complex, which is a coactivator for DNA-binding factors that activate transcription via RNA polymerase II (Sato et al., 2003 [PubMed 12584197]).[supplied by OMIM, Oct 2008],function:Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator

functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.,similarity:Belongs to the Mediator complex subunit 11 family.,subunit:Component of the Mediator complex, which is composed of MED1, MED4, MED6, MED7, MED8, MED9, MED10, MED11, MED12, MED13, MED13L, MED14, MED15, MED16, MED17, MED18, MED19, MED20, MED21, MED22, MED23, MED24, MED25, MED26, MED27, MED29, MED30, MED31, CCNC, CDK8 and CDC2L6/CDK11. The MED12, MED13, CCNC and CDK8 subunits form a distinct module termed the CDK8 module. Mediator containing the CDK8 module is less active than Mediator lacking this module in supporting transcriptional activation. Individual preparations of the Mediator complex lacking one or more distinct subunits have been variously termed ARC, CRSP, DRIP, PC2, SMCC and TRAP.,

## Research Area

## Image Data



Western blot analysis of lysates from SH-SY5Y cells, MED11 Rabbit Polyclonal Antibody was diluted at 1:1000, 4°over night