
Product Name: mGluR-4 Rabbit Polyclonal Antibody**Catalog #: APRab13861**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
Molecular Weight	100kDa

Antigen Information

Gene Name	GRM4
Alternative Names	GRM4; GPRC1D; MGLUR4; Metabotropic glutamate receptor 4; mGluR4
Gene ID	2914.0
SwissProt ID	Q14833
Immunogen	The antiserum was produced against synthesized peptide derived from human mGluR4. AA range:851-900

Background


glutamate metabotropic receptor 4(GRM4) Homo sapiens L-glutamate is the major excitatory neurotransmitter in the central

nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Several transcript variants encoding different isoforms have been found for this function: Receptor for glutamate. The activity of this receptor is mediated by a G-protein that inhibits adenylate cyclase activity., similarity: Belongs to the G-protein coupled receptor 3 family., subunit: Interacts with PICK1., tissue specificity: Strongly expressed in the cerebellum. Expressed at low levels in hippocampus, hypothalamus and thalamus. No expression detected in liver.,


Research Area

Neuroactive ligand-receptor interaction; Taste transduction;


Image Data




Immunohistochemistry analysis of paraffin-embedded human brain tissue, using mGluR4 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from 293 cells, treated with Forskolin 40nM 30', using mGluR4 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using mGluR-4 Polyclonal Antibody diluted at 1: 500



Western Blot analysis of 293 cells using mGluR-4 Polyclonal Antibody diluted at 1: 500