

Product Name: MKP-4 Rabbit Polyclonal Antibody

Catalog #: APRab13935

For research use only.

Summary

Description Rabbit polyclonal Antibody

Host Rabbit

Application WB,IHC,ICC/IF,ELISA
Reactivity Human,Mouse,Rat
Conjugation Unconjugated
Modification Unmodified

Isotype IgG

ClonalityPolyclonalFormLiquidConcentration1mg/ml

Storage Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

Shipping Ice bags

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer**

preservative N.

Purification Affinity purification

Application

Dilution Ratio WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000

Molecular Weight 42kDa

Antigen Information

Alternative Names

Gene Name DUSP9

DUSP9; MKP4; Dual specificity protein phosphatase 9; Mitogen-activated protein kinase

phosphatase 4; MAP kinase phosphatase 4; MKP-4

 Gene ID
 1852.0

 SwissProt ID
 Q99956

The antiserum was produced against synthesized peptide derived from human DUSP9. AA Immunogen

range:151-200

Background

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




The protein encoded by this gene is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which is associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product shows selectivity for members of the ERK family of MAP kinases and is localized to the cytoplasm and nucleus. Aberrant expression of this gene is associated with type 2 diabetes and cancer progreatalytic activity: A phosphoprotein + H(2)O = a protein + phosphate., catalytic activity: Protein tyrosine phosphate + H(2)O = protein tyrosine + phosphate., function: Inactivates MAP kinases. Has a specificity for the ERK family., similarity: Belongs to the protein-tyrosine phosphatase family. Non-receptor class dual specificity subfamily, similarity: Contains 1 rhodanese domain., similarity: Contains 1 tyrosine-protein phosphatase domain.,


Research Area

MAPK ERK Growth; MAPK G Protein;


Image Data



Immunofluorescence analysis of HeLa cells, using DUSP9 Antibody. The picture on the right is blocked with the synthesized peptide.




Immunohistochemistry analysis of paraffin-embedded human liver carcinoma tissue, using DUSP9 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa cells, using DUSP9 Antibody. The lane on the right is blocked with the synthesized peptide.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Western Blot analysis of COLO cells using MKP-4 Polyclonal Antibody diluted at 1: 1000

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838