

---

**Product Name: M-RIP Rabbit Polyclonal Antibody****Catalog #: APRab14095**

For research use only.

**Summary**

<b>Description</b>	Rabbit polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC,ELISA
<b>Reactivity</b>	Human,Mouse,Rat
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Concentration</b>	1mg/ml
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

**Application**

<b>Dilution Ratio</b>	WB 1:500-1:2000,IHC 1:50-1:300,ELISA 1:2000-1:20000
<b>Molecular Weight</b>	115kDa

**Antigen Information**

<b>Gene Name</b>	MPRIP
<b>Alternative Names</b>	MPRIP; KIAA0864; MRIP; RHOIP3; Myosin phosphatase Rho-interacting protein; M-RIP; Rho-interacting protein 3; RIP3; p116Rip
<b>Gene ID</b>	23164/23164
<b>SwissProt ID</b>	Q6WCQ1
<b>Immunogen</b>	Synthesized peptide derived from the Internal region of human M-RIP.


**Background**

function:Targets myosin phosphatase to the actin cytoskeleton. Required for the regulation of the actin cytoskeleton by RhoA


and ROCK1. Depletion leads to an increased number of stress fibers in smooth muscle cells through stabilization of actin fibers by phosphorylated myosin. Overexpression of MRIP as well as its F-actin-binding region leads to disassembly of stress fibers in neuronal cells.,similarity:Contains 1 PH domain.,similarity:Contains 2 PH domains.,subcellular location:Colocalizes with F-actin.,subunit:Binds F-actin through its N-terminus (By similarity). Binds RHOA, PPP1R12A/MBS and PPP1R12C/MBS85 through adjacent coiled coil domains.,function:Targets myosin phosphatase to the actin cytoskeleton. Required for the regulation of the actin cytoskeleton by RhoA and ROCK1. Depletion leads to an increased number of stress fibers in smooth muscle cells through stabilization of actin fibers by phosphorylated myosin. Overexpression of MRIP as well as its F-actin-binding region leads to disassembly of stress fibers in neuronal cells.,similarity:Contains 1 PH domain.,similarity:Contains 2 PH domains.,subcellular location:Colocalizes with F-actin.,subunit:Binds F-actin through its N-terminus (By similarity). Binds RHOA, PPP1R12A/MBS and PPP1R12C/MBS85 through adjacent coiled coil domains.,

## Research Area

### Image Data



Western Blot analysis of various cells using M-RIP Polyclonal Antibody



Western Blot analysis of 293 cells using M-RIP Polyclonal Antibody