

Product Name: NEIL2 Rabbit Polyclonal Antibody**Catalog #: APRab14546**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ELISA 1:5000-1:20000
Molecular Weight	36kDa

Antigen Information

Gene Name	NEIL2
Alternative Names	
Gene ID	252969.0
SwissProt ID	Q969S2
Immunogen	Synthesized peptide derived from part region of human protein

Background


NEIL2 belongs to a class of DNA glycosylases homologous to the bacterial Fpg/Nei family. These glycosylases initiate the first step in base excision repair by cleaving bases damaged by reactive oxygen species and introducing a DNA strand break via the associated lyase reaction (Bandaru et al., 2002 [PubMed 12509226])[supplied by OMIM, Mar 2008],catalytic activity:Removes

damaged bases from DNA, leaving an abasic site.,catalytic activity:The C-O-P bond 3' to the apurinic or apyrimidinic site in DNA is broken by a beta-elimination reaction, leaving a 3'-terminal unsaturated sugar and a product with a terminal 5'-phosphate.,domain:The zinc-finger domain is important for DNA binding.,enzyme regulation:Acetylation of Lys-50 leads to loss of DNA nicking activity. Acetylation of Lys-154 has no effect.,function:Involved in base excision repair of DNA damaged by oxidation or by mutagenic agents. Has DNA glycosylase activity towards 5-hydroxyuracil and other oxidized derivatives of cytosine with a preference for mismatched double stranded DNA (DNA bubbles). Has low or no DNA glycosylase activity towards thymine glycol, 2-hydroxyadenine, hypoxanthine and 8-oxoguanine. Has AP (apurinic/apyrimidinic) lyase activity and introduces nicks in the DNA strand. Cleaves the DNA backbone by beta-delta elimination to generate a single-strand break at the site of the removed base with both 3'- and 5'-phosphates.,similarity:Belongs to the FPG family.,similarity:Contains 1 FPG-type zinc finger.,subunit:Binds EP300.,tissue specificity:Detected in testis, skeletal muscle, heart, brain, placenta, lung, pancreas, kidney and liver.,

Research Area

Base excision repair;

Image Data



Western blot analysis of lysates from DU145 cells, NEIL2 Rabbit Polyclonal Antibody was diluted at 1:1000, 4° over night