

Product Name: Neuron Navigator 1 Rabbit Polyclonal Antibody**Catalog #: APRab14614**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	IHC, ICC/IF, ELISA
Reactivity	Human, Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:20000-1:40000
Molecular Weight	200kDa

Antigen Information

Gene Name	NAV1
Alternative Names	NAV1; KIAA1151; KIAA1213; POMFIL3; STEERIN1; Neuron navigator 1; Pore membrane and/or filament-interacting-like protein 3; Steerin-1; Unc-53 homolog 1; unc53H1
Gene ID	89796.0
SwissProt ID	Q8NEY1
Immunogen	Synthesized peptide derived from Neuron Navigator 1 . at AA range: 510-590

Background


This gene belongs to the neuron navigator family and is expressed predominantly in the nervous system. The encoded protein

contains coiled-coil domains and a conserved AAA domain characteristic for ATPases associated with a variety of cellular activities. This gene is similar to unc-53, a *Caenorhabditis elegans* gene involved in axon guidance. The exact function of this gene is not known, but it is thought to play a role in neuronal development and regeneration. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2009],developmental stage:Expressed in fetal brain and heart.,function:May be involved in neuronal migration.,similarity:Belongs to the Nav/unc-53 family.,subcellular location:Associates with a subset of microtubule plus ends. Enriched in neuronal growth cones.,subunit:Interacts with tubulin.,tissue specificity:Broadly expressed at low levels. Expressed at high levels in heart, skeletal muscle and placenta.,

Research Area

Growth and Development; Axonal Guidance Proteins; Neuroscience; Neurology process; Neurogenesis

Image Data



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200 (4° overnight) . 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min) .