

## **Product Name: NOXIN Rabbit Polyclonal Antibody**

Catalog #: APRab14820

For research use only.

### **Summary**

**Description** Rabbit polyclonal Antibody

Host Rabbit
Application WB,ELISA

Reactivity Human,Rat,Mouse
Conjugation Unconjugated
Modification Unmodified

**Isotype** IgG

ClonalityPolyclonalFormLiquidConcentration1mg/ml

Storage Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

**Shipping** Ice bags

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer** 

preservative N.

**Purification** Affinity purification

## **Application**

**Dilution Ratio** WB 1:500-1:2000,ELISA 1:5000-1:20000

Molecular Weight 85kDa

# **Antigen Information**

Gene Name NOXIN

Alternative Names NOXIN; C11orf82; Nitric oxide-inducible gene protein

 Gene ID
 220042.0

 SwissProt ID
 Q8IXT1

The antiserum was produced against synthesized peptide derived from the Internal region of Immunogen

human NOXIN. AA range:111-160

## **Background**

function: Acts as an anti-apoptotic factor and its absence increases cell death under normal and stress conditions. Can induce


Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838



cell cycle arrest in the G1 or early S phase and this activity is independent of TP53/p53.,subcellular location:Accumulates in the nucleus in response to stress.,function:Acts as an anti-apoptotic factor and its absence increases cell death under normal and stress conditions. Can induce cell cycle arrest in the G1 or early S phase and this activity is independent of TP53/p53.,subcellular location:Accumulates in the nucleus in response to stress.,

#### **Research Area**

## **Image Data**



Western Blot analysis of 823, 293T, AD293, Hela, HepG2 cells using NOXIN Polyclonal Antibody. Antibody was diluted at 1:2000. Secondary antibody was diluted at 1:20000