

Product Name: OAS2 Rabbit Polyclonal Antibody**Catalog #: APRab15078**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Molecular Weight	82kDa

Antigen Information

Gene Name	OAS2
Alternative Names	OAS2; 2'-5'-oligoadenylate synthase 2; (2-5')oligo(A) synthase 2; 2-5A synthase 2; p69 OAS / p71 OAS; p69OAS / p71OAS
Gene ID	4939.0
SwissProt ID	P29728
Immunogen	The antiserum was produced against synthesized peptide derived from the N-terminal region of human OAS2. AA range:61-110


Background

2'-5'-oligoadenylate synthetase 2(OAS2) Homo sapiens This gene encodes a member of the 2-5A synthetase family, essential proteins involved in the innate immune response to viral infection. The encoded protein is induced by interferons and uses adenosine triphosphate in 2'-specific nucleotidyl transfer reactions to synthesize 2',5'-oligoadenylates (2-5As). These molecules activate latent RNase L, which results in viral RNA degradation and the inhibition of viral replication. The three known members of this gene family are located in a cluster on chromosome 12. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq, Jul 2008],catalytic activity: Binds double-stranded RNA and polymerizes ATP into PPP(A2'P5'A)N oligomers, which activate the latent RNase L that, when activated, cleaves single-stranded RNAs.,function: May play a role in mediating resistance to virus infection, control of cell growth, differentiation, and apoptosis.,induction: By interferons.,similarity: Belongs to the 2-5A synthetase family.,subcellular location: Associated with different subcellular fractions such as mitochondrial, nuclear, and rough/smooth microsomal fractions.,subunit: Homodimer.,


Research Area

DNA / RNA; RNA Processing; Epigenetics and Nuclear Signaling; Chromatin Binding Proteins; DNA / RNA binding; Microbiology; Organism; Virus; RNA Virus; ssRNA positive strand virus; SARS Coronavirus; Immunology; Innate Immunity; Cytokines; Interferons


Image Data



Western Blot analysis of HeLa cells using OAS2 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-uterus, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-uterus, antibody was diluted at 1:100