

# **Product Name:** Olfactory receptor 10C1 Rabbit Polyclonal Antibody

Catalog #: APRab15132

For research use only.

## **Summary**

**Description** Rabbit polyclonal Antibody

Host Rabbit
Application WB,ELISA

Reactivity Human,Rat,Mouse
Conjugation Unconjugated
Modification Unmodified

**Isotype** IgG

ClonalityPolyclonalFormLiquidConcentration1mg/ml

**Storage** Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

**Shipping** Ice bags

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer** 

preservative N.

**Purification** Affinity purification

## **Application**

**Dilution Ratio** WB 1:500-1:2000,ELISA 1:20000-1:40000

Molecular Weight 34kDa

# **Antigen Information**

Gene Name OR10C1

**Alternative Names** 

**Gene ID** 442194.0

**SwissProt ID** Q96KK4/Q5SUN7

The antiserum was produced against synthesized peptide derived from human OR10C1. AA Immunogen

range:61-110

## **Background**

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a




smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. This olfactory receptor gene is a segregating pseudogene, where some individuals have an allele that encodes a functional olfactory receptor, while other individuals have an allele encoding afunction:Odorant receptor "polymorphism:A stop codon at position Gln-55 in the gene coding for this protein is responsible for functional diversity thus producing a pseudogene. The stop codon is more frequent in African-Americans than in non-Africans, similarity:Belongs to the G-protein coupled receptor 1 family.


### **Research Area**

Olfactory transduction;

#### **Image Data**



Western blot analysis of lysates from LOVO cells, using OR10C1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HeLa cells using OR10C1 antibody.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838