# **Product Name: Olfactory receptor 13C8 Rabbit**

Polyclonal Antibody Catalog #: APRab15160



# **Summary**

**Production Name** Olfactory receptor 13C8 Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

Host Rabbit
Application WB,ELISA

**Reactivity** Human, Rat, Mouse

### **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

Clonality Polyclonal Form Liquid

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type

preservative N.

**Purification** Affinity purification

### **Immunogen**

**Buffer** 

Gene Name OR13C8

Alternative Names OR13C8; Olfactory receptor 13C8

Gene ID 138802.0

Q8NGS7. The antiserum was produced against synthesized peptide derived from **SwissProt ID** 

human OR13C8. AA range:271-320

# **Application**

**Dilution Ratio** WB 1:500-1:2000, ELISA 1:5000.Not yet tested in other applications.

Molecular Weight 35kDa

 **Product Name: Olfactory receptor 13C8 Rabbit** 

Polyclonal Antibody Catalog #: APRab15160




# **Background**

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008], function:Odorant receptor ., similarity:Belongs to the G-protein coupled receptor 1 family.,

### Research Area

Olfactory transduction;

# **Image Data**



Western blot analysis of lysates from NIH/3T3 cells, using OR13C8 Antibody. The lane on the right is blocked with the synthesized peptide.

### Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838