

# **Product Name: Olfactory receptor 2AJ1 Rabbit Polyclonal Antibody**

Catalog #: APRab15179

For research use only.

#### **Summary**

**Description** Rabbit polyclonal Antibody

**Host** Rabbit

Application WB,ICC/IF,ELISA
Reactivity Human,Rat,Mouse
Conjugation Unconjugated
Modification Unmodified

**Isotype** IgG

ClonalityPolyclonalFormLiquidConcentration1mg/ml

**Storage** Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

**Shipping** Ice bags

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer** 

preservative N.

**Purification** Affinity purification

#### **Application**

**Dilution Ratio** WB 1:500-1:2000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:10000

Molecular Weight 42kDa

### **Antigen Information**

Gene Name OR2AJ1

Alternative Names OR2AJ1; OR2AJ1P; Olfactory receptor 2AJ1

 Gene ID
 127608.0

 SwissProt ID
 Q8NGZ0

The antiserum was produced against synthesized peptide derived from human OR2AJ1. AA Immunogen

range:201-250

#### **Background**

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a


Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,

#### **Research Area**


## **Image Data**



Western blot analysis of lysates from Jurkat and HUVEC cells, using OR2AJ1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using OR2AJ1 antibody.



Western Blot analysis of Jurkat cells using Olfactory receptor 2AJ1 Polyclonal Antibody

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

