

Product Name: Olfactory receptor 2T2/35 Rabbit Polyclonal Antibody Catalog #: APRab15205

For research use only.

Summary

Description Rabbit polyclonal Antibody

Host Rabbit

Application WB,ICC/IF,ELISA
Reactivity Human,Rat,Mouse
Conjugation Unconjugated
Modification Unmodified

Isotype IgG

ClonalityPolyclonalFormLiquidConcentration1mg/ml

Storage Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

Shipping Ice bags

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer**

preservative N.

Purification Affinity purification

Application

Dilution Ratio WB 1:500-1:2000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000

Molecular Weight 33kDa

Antigen Information

Alternative Names

Gene Name OR2T2/OR2T35

OR2T2; OR2T2P; Olfactory receptor 2T2; Olfactory receptor OR1-43; OR2T35; Olfactory

receptor 2T35; Olfactory receptor OR1-66

 Gene ID
 401992/403244

 SwissProt ID
 Q6IF00/Q8NGX2

The antiserum was produced against synthesized peptide derived from human OR2T2/2T35.

AA range:61-110

Background




olfactory receptor family 2 subfamily T member 2(OR2T2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008], function:Odorant receptor ., similarity:Belongs to the G-protein coupled receptor 1 family.,

Research Area


Olfactory transduction;

Image Data



Immunofluorescence analysis of A549 cells, using OR2T2/2T35 Antibody. The picture on the right is blocked with the synthesized peptide.


Western blot analysis of lysates from HepG2 cells, using OR2T2/2T35 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HeLa cells using OR2T2/35 antibody.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Western Blot analysis of various cells using Olfactory receptor 2T2/35 Polyclonal Antibody

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838