

# **Product Name: Olfactory receptor 2W3 Rabbit Polyclonal Antibody**

Catalog #: APRab15208

For research use only.

#### **Summary**

**Description** Rabbit polyclonal Antibody

Host Rabbit
Application WB,ELISA

Reactivity Human,Rat,Mouse
Conjugation Unconjugated
Modification Unmodified

**Isotype** IgG

ClonalityPolyclonalFormLiquidConcentration1mg/ml

**Storage** Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

**Shipping** Ice bags

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer** 

preservative N.

**Purification** Affinity purification

### **Application**

**Dilution Ratio** WB 1:500-1:2000,ELISA 1:5000-1:20000

Molecular Weight 35kDa

## **Antigen Information**

**Alternative Names** 

Gene Name OR2W3

OR2W3; OR2W3P; OR2W8P; Olfactory receptor 2W3; Olfactory receptor 2W8; Olfactory

receptor OR1-49

 Gene ID
 343171.0

 SwissProt ID
 O7Z3T1

The antiserum was produced against synthesized peptide derived from human OR2W3. AA Immunogen

range:141-190

## **Background**




olfactory receptor family 2 subfamily W member 3(OR2W3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,


#### **Research Area**

Olfactory transduction;

#### **Image Data**



Western blot analysis of lysates from NIH/3T3 cells, using OR2W3 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of 3T3 cells using Olfactory receptor 2W3 Polyclonal Antibody

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838