
Product Name: Olfactory receptor O13C2/9 Rabbit Polyclonal Antibody**Catalog #: APRab15342**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ICC/IF,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
Molecular Weight	36kDa

Antigen Information

Gene Name	OR13C2/OR13C9
Alternative Names	OR13C2; Olfactory receptor 13C2; Olfactory receptor OR9-12; OR13C9; Olfactory receptor 13C9; Olfactory receptor OR9-13
Gene ID	392376/286362
SwissProt ID	Q8NGS9/Q8NGT0
Immunogen	The antiserum was produced against synthesized peptide derived from human OR13C2/13C9. AA range:232-281


Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],function:Odorant receptor „similarity:Belongs to the G-protein coupled receptor 1 family,.


Research Area

Olfactory transduction;

Image Data



Western blot analysis of lysates from Jurkat cells, using OR13C2/13C9 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from K562 cells using OR13C2/13C9 antibody.