
Product Name: POLD4 Rabbit Polyclonal Antibody**Catalog #: APRab16348**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:10000

Molecular Weight

Antigen Information

Gene Name	POLD4 POLDS
Alternative Names	DNA polymerase delta subunit 4 (DNA polymerase delta subunit p12)
Gene ID	57804.0
SwissProt ID	Q9HCU8
Immunogen	Synthesized peptide derived from human POLD4 AA range: 43-93

Background


This gene encodes the smallest subunit of DNA polymerase delta. DNA polymerase delta possesses both polymerase and 3' to 5' exonuclease activity and plays a critical role in DNA replication and repair. The encoded protein enhances the activity of DNA

polymerase delta and plays a role in fork repair and stabilization through interactions with the DNA helicase Bloom syndrome protein. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Mar 2012],function:Required for optimal DNA polymerase delta activity. May contribute to PCNA-dependent activity of DNA polymerase delta.,similarity:Belongs to the DNA polymerase delta subunit 4 family.,subunit:Heterotetramer composed of subunits of 125 kDa, 50 kDa, 66 kDa and 12 kDa. Interacts with WRNIP1. Interacts with POLD4, POLD2, POLD1 and PCNA.,

Research Area

Epigenetics and Nuclear Signaling; DNA / RNA; DNA Synthesis; DNA Polymerases

Image Data



Immunohistochemical analysis of paraffin-embedded human Small intestinal stromal tumor. 1, Tris-EDTA,pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200 (4° overnight.3,Secondary antibody was diluted at 1:200 (room temperature, 45min) .