

Product Name: PP32R1 Rabbit Polyclonal Antibody

Catalog #: APRab16402

For research use only.

Summary

Description Rabbit polyclonal Antibody

Host Rabbit
Application WB,IHC

Reactivity Human,Rat,Mouse
Conjugation Unconjugated
Modification Unmodified

Isotype IgG

ClonalityPolyclonalFormLiquidConcentration1mg/ml

Storage Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

Shipping Ice bags

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer**

preservative N.

Purification Affinity purification

Application

Dilution Ratio WB 1:500-1:2000,IHC 1:50-1:300

Molecular Weight 26kDa

Antigen Information

Alternative Names

Gene Name ANP32C

ANP32C; PP32R1; Acidic leucine-rich nuclear phosphoprotein 32 family member C;

Phosphoprotein 32-related protein 1; Tumorigenic protein pp32r1

 Gene ID
 23520.0

 SwissProt ID
 043423

The antiserum was produced against synthesized peptide derived from human ANP32C. AA Immunogen

range:121-170

Background




Phosphoprotein 32 (PP32) is a tumor suppressor that can inhibit several types of cancers, including prostate and breast cancers. The protein encoded by this gene is one of at least two proteins that are similar in amino acid sequence to PP32 and are part of the same acidic nuclear phosphoprotein gene family. However, unlike PP32, the encoded protein is tumorigenic. The tumor suppressor function of PP32 has been localized to a 25 amino acid region that is divergent between PP32 and the protein encoded by this gene. This gene does not contain introns. [provided by RefSeq, Jul 2008],caution:The His-140 polymorphism is uncertain. It has been found in prostatic cell line without corresponding normal tissue sample. Cells transfected with this variant expressed at least twice as much proteins than native cells.,similarity:Belongs to the ANP32 family.,similarity:Contains 4 LRR (leucine-rich) repeats.,tissue specificity:Expressed in activated stem cells, such as mobilized CD34+ cells and cord blood CD34+ cells, but not in resting bone marrow CD34+ cells. Expressed in a variety of neoplastic cell lines, mainly in prostatic adenocarcinoma cell lines. Not expressed in normal protastic tissue.,


Research Area

Cancer; Oncoproteins/suppressors; Oncoproteins

Image Data



Western blot analysis of lysates from HUVEC cells, using ANP32C Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using PP32R1 Polyclonal Antibody diluted at 1 : 1000

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838