

Product Name: PPP1R3A Rabbit Polyclonal Antibody**Catalog #: APRab16428**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:50-1:300
Molecular Weight	140kDa

Antigen Information

Gene Name	PPP1R3A
Alternative Names	PPP1R3A; PP1G; Protein phosphatase 1 regulatory subunit 3A; Protein phosphatase 1 glycogen-associated regulatory subunit; Protein phosphatase type-1 glycogen targeting subunit; RG1
Gene ID	5506.0
SwissProt ID	Q16821
Immunogen	The antiserum was produced against synthesized peptide derived from human PPP1R3A. AA range:647-696


Background

The glycogen-associated form of protein phosphatase-1 (PP1) derived from skeletal muscle is a heterodimer composed of a 37-kD catalytic subunit and a 124-kD targeting and regulatory subunit. This gene encodes the regulatory subunit which binds to muscle glycogen with high affinity, thereby enhancing dephosphorylation of glycogen-bound substrates for PP1 such as glycogen synthase and glycogen phosphorylase kinase. [provided by RefSeq, Jul 2008],disease:Defects in PPP1R3A are a cause of insulin resistance (Ins resistance),disease:Defects in PPP1R3A are a cause of susceptibility to noninsulin-dependent diabetes mellitus (NIDDM) [MIM:125853]; also known as diabetes mellitus type II. NIDDM is characterized by an autosomal dominant mode of inheritance, onset during adulthood and insulin resistance.,domain:The CBM21 domain is known to be involved in the localization to glycogen and is characteristic of some regulatory subunit of phosphatase complexes.,function:Seems to act as a glycogen-targeting subunit for PP1. PP1 is essential for cell division, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. Plays an important role in glycogen synthesis but is not essential for insulin activation of glycogen synthase.,PTM:Phosphorylation at Ser-46 by ISPK stimulates the dephosphorylation of glycogen synthase and phosphorylase kinase.,similarity:Contains 1 CBM21 (carbohydrate binding type-21) domain.,subunit:Interacts with PPP1CC catalytic subunit of PP1, and associates with glycogen.,tissue specificity:Skeletal muscle and heart.,


Research Area

Insulin_Receptor;

Image Data



Western blot analysis of lysates from MCF-7 and COLO cells, using PPP1R3A Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using PPP1R3A antibody.

