

---

**Product Name: PRP6 Rabbit Polyclonal Antibody****Catalog #: APRab16542**

For research use only.

**Summary**

<b>Description</b>	Rabbit polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC,ICC/IF,ELISA
<b>Reactivity</b>	Human,Mouse,Rat
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Concentration</b>	1mg/ml
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

**Application**

<b>Dilution Ratio</b>	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
<b>Molecular Weight</b>	110kDa

**Antigen Information**

<b>Gene Name</b>	PRPF6 PRPF6; C20orf14; Pre-mRNA-processing factor 6; Androgen receptor N-terminal domain-
<b>Alternative Names</b>	transactivating protein 1; ANT-1; PRP6 homolog; U5 snRNP-associated 102 kDa protein; U5-102 kDa protein
<b>Gene ID</b>	24148.0
<b>SwissProt ID</b>	O94906
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human PRPF6. AA range:747-796


## Background

The protein encoded by this gene appears to be involved in pre-mRNA splicing, possibly acting as a bridging factor between U5 and U4/U6 snRNPs in formation of the spliceosome. The encoded protein also can bind androgen receptor, providing a link between transcriptional activation and splicing. [provided by RefSeq, Jul 2008],function:Involved in pre-mRNA splicing. May act in the tri-snRNP complex as a bridging factor between U5 and U4/U6 snRNPs in the late step of spliceosome assembly. May be necessary for tri-snRNP formation.,similarity:Contains 9 HAT repeats.,subunit:Associates with the U5 snRNP particle (containing the 40 kDa, 52 kDa, 116 kDa, 200 kDa and 220 kDa proteins) probably via interaction(s) with the 220 kDa and/or other proteins. Associates with U4/U6 snRNP particle (containing the 15.5 kDa, 20 kDa/60kDa/90kDa heteromer, LSm proteins LSm2-8, 61 kDa and Sm proteins). Interacts with ARAF1. Identified in the spliceosome C complex, at least composed of AQR, ASCC3L1, C19orf29, CDC40, CDC5L, CRNKL1, DDX23, DDX41, DDX48, DDX5, DGCR14, DHX35, DHX38, DHX8, EFTUD2, FRG1, GPATC1, HNRPA1, HNRPA2B1, HNRPA3, HNRPC, HNRPF, HNRPH1, HNRPK, HNRPM, HNRPR, HNRPU, KIAA1160, KIAA1604, LSM2, LSM3, MAGOH, MORG1, PABPC1, PLRG1, PNN, PPIE, PPIL1, PPIL3, PPWD1, PRPF19, PRPF4B, PRPF6, PRPF8, RALY, RBM22, RBM8A, RBMX, SART1, SF3A1, SF3A2, SF3A3, SF3B1, SF3B2, SF3B3, SFRS1, SKIV2L2, SNRPA1, SNRPB, SNRPB2, SNRPD1, SNRPD2, SNRPD3, SNRPE, SNRPF, SNRPG, SNW1, SRRM1, SRRM2, SYF2, SYNCRIP, TFIP11, THOC4, U2AF1, WDR57, XAB2 and ZCCHC8.,

## Research Area

Spliceosome;

## Image Data



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using PRPF6 Antibody. The picture on the right is blocked with the synthesized peptide.