
Product Name: RNase III Drosha Rabbit Polyclonal Antibody**Catalog #: APRab17277**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
Molecular Weight	160kDa

Antigen Information

Gene Name	DROSHA
Alternative Names	DROSHA; RN3; RNASE3L; RNASEN; Ribonuclease 3; Protein Drosha; Ribonuclease III; RNase III; p241
Gene ID	29102.0
SwissProt ID	Q9NRR4
Immunogen	The antiserum was produced against synthesized peptide derived from human RNase III Drosha. AA range:774-823


Background

drosha ribonuclease III(DROSHA) Homo sapiens This gene encodes a ribonuclease (RNase) III double-stranded RNA-specific ribonuclease and subunit of the microprocessor protein complex, which catalyzes the initial processing step of microRNA (miRNA) synthesis. The encoded protein cleaves the stem loop structure from the primary microRNA (pri-miRNA) in the nucleus, yielding the precursor miRNA (pre-miRNA), which is then exported to the cytoplasm for further processing. In a human cell line lacking a functional copy of this gene, canonical miRNA synthesis is reduced. Somatic mutations in this gene have been observed in human patients with kidney cancer. [provided by RefSeq, Sep 2016],catalytic activity:Endonucleolytic cleavage to 5'-phosphomonoester.,cofactor:Magnesium or manganese.,function:Executes the initial step of microRNA (miRNA) processing in the nucleus, that is cleavage of pri-miRNA to release pre-miRNA. Involved in pre-rRNA processing. Cleaves double-strand RNA and does not cleave single-strand RNA.,online information:The dark side of RNA -Issue 87 of October 2007,similarity:Contains 1 DRBM (double-stranded RNA-binding) domain.,similarity:Contains 2 RNase III domains.,subcellular location:A fraction is translocated to the nucleolus during the S phase of the cell cycle.,subunit:Interacts with Sp1.,tissue specificity:Ubiquitous.,

Research Area


Epigenetics and Nuclear Signaling; DNA / RNA; RNA Processing RNAi; Dicer; Nuclear Signaling Pathways; Nuclear Receptors; Nuclear Pore Complex

Image Data




Western blot analysis of lysates from brain tissue, using RNase III Drosha antibody.


Western blot analysis of SH-SY5Y K562 HELA using RNase III Drosha antibody. Antibody was diluted at 1:1000. Secondary antibody was diluted at 1:20000.



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200