

Product Name: SR-3D Rabbit Polyclonal Antibody**Catalog #: APRab18251**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ICC/IF,ELISA
Reactivity	Human
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000
Molecular Weight	50kDa

Antigen Information

Gene Name	HTR3D
Alternative Names	HTR3D; 5-hydroxytryptamine receptor 3D; 5-HT3-D; 5-HT3D; Serotonin receptor 3D
Gene ID	200909.0
SwissProt ID	Q70Z44
Immunogen	Synthesized peptide derived from SR-3D . at AA range: 10-90


Background

The protein encoded this gene belongs to the ligand-gated ion channel receptor superfamily. This gene encodes subunit D of the type 3 receptor for 5-hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a mitogen

and a hormone. This hormone has been linked to neuropsychiatric disorders, including anxiety, depression, and migraine. Serotonin receptors causes fast and depolarizing responses in neurons following activation. The genes encoding subunits C, D and E of this type 3 receptor form a cluster on chromosome 3. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2009],function:This is one of the several different receptors for 5-hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone, and a mitogen. This receptor is a ligand-gated ion channel, which when activated causes fast, depolarizing responses. It is a cation-specific, but otherwise relatively nonselective, ion channel.,similarity:Belongs to the ligand-gated ionic channel (TC 1.A.9) family.,subcellular location:Presumably retained within the endoplasmic reticulum unless complexed with HTR3A.,subunit:Forms a pentaheteromeric complex with HTR3A, homomeric complex being not functional.,tissue specificity:Expressed in liver, as well as fetal and adult colon and kidney.,

Research Area

Image Data



Western Blot analysis of HT-29 cells using SR-3D Polyclonal Antibody