
Product Name: SV2C Rabbit Polyclonal Antibody**Catalog #: APRab18464**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Molecular Weight	82kDa

Antigen Information

Gene Name	SV2C
Alternative Names	SV2C; KIAA1054; Synaptic vesicle glycoprotein 2C
Gene ID	22987.0
SwissProt ID	Q496J9
Immunogen	Synthesized peptide derived from the Internal region of human SV2C. AA range 330-390

Background


function:Plays a role in the control of regulated secretion in neural and endocrine cells, enhancing selectively low-frequency neurotransmission. Positively regulates vesicle fusion by maintaining the readily releasable pool of secretory vesicles.,PTM:N-

glycosylated.,similarity:Belongs to the major facilitator superfamily.,subcellular location:Enriched in small synaptic vesicles and adrenal microsomes, not present in chromaffin granules. Associated with both insulin granules and synaptic-like microvesicles in insulin-secreting cells of the pancreas.,subunit:Interacts with SYT1 in a calcium-dependent manner.,function:Plays a role in the control of regulated secretion in neural and endocrine cells, enhancing selectively low-frequency neurotransmission. Positively regulates vesicle fusion by maintaining the readily releasable pool of secretory vesicles.,PTM:N-glycosylated.,similarity:Belongs to the major facilitator superfamily.,subcellular location:Enriched in small synaptic vesicles and adrenal microsomes, not present in chromaffin granules. Associated with both insulin granules and synaptic-like microvesicles in insulin-secreting cells of the pancreas.,subunit:Interacts with SYT1 in a calcium-dependent manner.,


Research Area

ECM-receptor interaction;

Image Data



Immunohistochemical analysis of paraffin-embedded rat-brain, antibody was diluted at 1:100



Western Blot analysis of extracts from Jurkat cells, using SV2C Polyclonal Antibody. Secondary antibody was diluted at 1:20000