

Product Name: TAAR5 Rabbit Polyclonal Antibody

Catalog #: APRab18590

For research use only.

Summary

Description Rabbit polyclonal Antibody

Host Rabbit
Application WB,ELISA

Reactivity Human,Rat,Mouse
Conjugation Unconjugated
Modification Unmodified

Isotype IgG

ClonalityPolyclonalFormLiquidConcentration1mg/ml

Storage Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

Shipping Ice bags

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer**

preservative N.

Purification Affinity purification

Application

Dilution Ratio WB 1:500-1:2000,ELISA 1:10000-1:20000

Molecular Weight 38kDa

Antigen Information

Alternative Names

Gene Name TAAR5

TAAR5; PNR; Trace amine-associated receptor 5; TaR-5; Trace amine receptor 5; Putative

neurotransmitter receptor

 Gene ID
 9038.0

 SwissProt ID
 O14804

The antiserum was produced against synthesized peptide derived from human TAAR5. AA **Immunogen**

range:288-337

Background




function:Orphan receptor. Ligands are likely small molecules, either sharing some similarities with trace amine as, e.g. derivatives of indolamines (such as 5-methoxytryptamine) or of phenylethylamines (such as phenylethanolamine) or being any kind of metabolite of amino acids or biogenic amine neurotransmitters., similarity:Belongs to the G-protein coupled receptor 1 family., tissue specificity:Expressed almost exclusively in skeletal muscle and selected areas of the brain, such amygdala, hippocampus, caudate nucleus, thalamus and hypothalamus. Weak expression is also find in substantia nigra., function:Orphan receptor. Ligands are likely small molecules, either sharing some similarities with trace amine as, e.g. derivatives of indolamines (such as 5-methoxytryptamine) or of phenylethylamines (such as phenylethanolamine) or being any kind of metabolite of amino acids or biogenic amine neurotransmitters., similarity:Belongs to the G-protein coupled receptor 1 family., tissue specificity:Expressed almost exclusively in skeletal muscle and selected areas of the brain, such amygdala, hippocampus, caudate nucleus, thalamus and hypothalamus. Weak expression is also find in substantia nigra.,

Research Area

Neuroactive ligand-receptor interaction;

Image Data



34-

26-

19-

Western blot analysis of TAAR5 Antibody. The lane on the right is blocked with the TAAR5 peptide.

Western blot analysis of the lysates from HUVECcells using TAAR5 antibody.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838