
Product Name: TALK-2 Rabbit Polyclonal Antibody**Catalog #: APRab18639**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ELISA 1:10000-1:20000
Molecular Weight	37-42kDa

Antigen Information


Gene Name	KCNK17 KCNK17; TALK2; TASK4; Potassium channel subfamily K member 17; 2P domain potassium channel
Alternative Names	Talk-2; Acid-sensitive potassium channel protein TASK-4; TWIK-related acid-sensitive K(+) channel 4; TWIK-related alkaline pH-activated K(+) channel
Gene ID	89822.0
SwissProt ID	Q96T54
Immunogen	The antiserum was produced against synthesized peptide derived from human KCNK17. AA range:271-320

Background

potassium two pore domain channel subfamily K member 17(KCNK17) Homo sapiens The protein encoded by this gene belongs to the family of potassium channel proteins containing two pore-forming P domains. This channel is an open rectifier which primarily passes outward current under physiological K⁺ concentrations. This gene is activated at alkaline pH. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2008],function:Outward rectifying potassium channel. Produces rapidly activating and non-inactivating outward rectifier K(+) currents.,miscellaneous:Inhibited by Ba(2+), quinidine, chloroform and halothane. Activated at alkaline pH. Activated by quinine and isoflurane.,similarity:Belongs to the two pore domain potassium channel (TC 1.A.1.8) family.,subunit:Homodimer .,

Research Area

Image Data



Western blot analysis of lysates from Jurkat cells, using KCNK17 Antibody. The lane on the right is blocked with the synthesized peptide.