

**Product Name: TCAM2 Rabbit Polyclonal Antibody****Catalog #: APRab18719**

For research use only.

**Summary**

<b>Description</b>	Rabbit polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,ELISA
<b>Reactivity</b>	Human,Mouse
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Concentration</b>	1mg/ml
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

**Application**

<b>Dilution Ratio</b>	WB 1:500-1:2000,ELISA 1:5000-1:20000
<b>Molecular Weight</b>	25kDa

**Antigen Information**

<b>Gene Name</b>	TICAM2 TIRAP3 TIRP TRAM
<b>Alternative Names</b>	
<b>Gene ID</b>	100302736.0
<b>SwissProt ID</b>	Q86XR7
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein

**Background**

TIRP is a Toll/interleukin-1 receptor (IL1R; MIM 147810) (TIR) domain-containing adaptor protein involved in Toll receptor signaling (see TLR4; MIM 603030).[supplied by OMIM, Apr 2004],domain:The TIR domain mediates the interaction with TRAF6,.function:Functions in LPS-TLR4 signaling to regulate the MYD88-independent pathway during the innate immune

response to LPS. Also involved in IL1-triggered NF-kappa-B activation, functioning upstream of IRAK1, IRAK2, TRAF6, and IKKB. Physically bridges TLR4 and TICAM1 and functionally transmits LPS-TLR4 signal to TICAM1.,PTM:Myristoylated. Required for membrane association which is critical for its ability to initiate efficient signaling.,PTM:Phosphorylated by PKCE in response to LPS. Phosphorylation is essential for its function. It is depleted from the membrane upon phosphorylation.,similarity:Belongs to the EMP24/GP25L family.,similarity:Contains 1 GOLD domain.,similarity:Contains 1 TIR domain.,subcellular location:Localized to the plasma membrane as a result of myristoylation. Phosphorylation on Ser-16 leads to its depletion from the membrane.,subunit:Homodimer. Interacts with TLR4, TICAM1, IRF3 and IRF7 in response to LPS. Interacts with IL1R1, IL1RAP, IRAK2, IRAK3 and TRAF6. Interacts with protein kinase-inactive mutants of IRAK1 and IRAK4.,tissue specificity:Expressed in spleen, prostate, testis, uterus, small intestine, colon, peripheral blood leukocytes, heart, placenta, lung, liver, skeletal muscle, and pancreas.,

## Research Area

Toll\_Like;

## Image Data

