

Product Name: TNNI3K Rabbit Polyclonal Antibody**Catalog #: APRab19105**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000

Molecular Weight

Antigen Information

Gene Name	TNNI3K
Alternative Names	TNNI3K; CARK; Serine/threonine-protein kinase TNNI3K; Cardiac ankyrin repeat kinase; Cardiac troponin I-interacting kinase; TNNI3-interacting kinase
Gene ID	51086.0
SwissProt ID	Q59H18
Immunogen	The antiserum was produced against synthesized peptide derived from human TNNI3K. AA range:301-350


Background

This gene encodes a protein that belongs to the MAP kinase kinase kinase (MAPKKK) family of protein kinases. The protein contains ankyrin repeat, protein kinase and serine-rich domains and is thought to play a role in cardiac physiology. [provided by RefSeq, Sep 2012], catalytic activity: ATP + a protein = ADP + a phosphoprotein., catalytic activity: GTP + beta-L-fucose 1-phosphate = diphosphate + GDP-L-fucose., cofactor: Magnesium., function: Catalyzes the formation of GDP-L-fucose from GTP and L-fucose-1-phosphate. Functions as a salvage pathway to reutilize L-fucose arising from the turnover of glycoproteins and glycolipids., function: May play a role in cardiac physiology., PTM: Autophosphorylated., similarity: Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family. MAP kinase kinase kinase subfamily., similarity: Contains 1 protein kinase domain., similarity: Contains 10 ANK repeats., subcellular location: Expressed at lower levels in the cytoplasm., subunit: Interacts with TNNI3, ACTC, ACTA1, MYBPC3, AIP, BABP3 and HADHB., tissue specificity: Expressed in many tissues., tissue specificity: Highly expressed in both adult and fetal heart.,


Research Area

Fructose and mannose metabolism; Amino sugar and nucleotide sugar metabolism;

Image Data



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using TNNI3K Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of Mouse-kidney lysis using TNNI3K antibody. Antibody was diluted at 1:500. Secondary antibody was diluted at 1:20000