

## Summary

|                        |                                 |
|------------------------|---------------------------------|
| <b>Production Name</b> | Ubr1 Rabbit Polyclonal Antibody |
| <b>Description</b>     | Rabbit Polyclonal Antibody      |
| <b>Host</b>            | Rabbit                          |
| <b>Application</b>     | WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA |
| <b>Reactivity</b>      | Human,Mouse                     |

## Performance

|                     |                                                                                                   |
|---------------------|---------------------------------------------------------------------------------------------------|
| <b>Conjugation</b>  | Unconjugated                                                                                      |
| <b>Modification</b> | Unmodified                                                                                        |
| <b>Isotype</b>      | IgG                                                                                               |
| <b>Clonality</b>    | Polyclonal                                                                                        |
| <b>Form</b>         | Liquid                                                                                            |
| <b>Storage</b>      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.          |
| <b>Buffer</b>       | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| <b>Purification</b> | Affinity purification                                                                             |

## Immunogen

|                          |                                                                                                                                |
|--------------------------|--------------------------------------------------------------------------------------------------------------------------------|
| <b>Gene Name</b>         | UBR1                                                                                                                           |
| <b>Alternative Names</b> | UBR1; E3 ubiquitin-protein ligase UBR1; N-recognin-1; Ubiquitin-protein ligase E3-alpha-1; Ubiquitin-protein ligase E3-alpha-I |
| <b>Gene ID</b>           | 197131.0                                                                                                                       |
| <b>SwissProt ID</b>      | Q8IWW7.The antiserum was produced against synthesized peptide derived from human UBR1. AA range:821-870                        |

## Application


|                         |                                                                              |
|-------------------------|------------------------------------------------------------------------------|
| <b>Dilution Ratio</b>   | WB 1:500-1:2000, IHC-P 1:100-1:300, ELISA 1:20000, IF-P/IF-F/ICC/IF 1:50-200 |
| <b>Molecular Weight</b> | 200kDa                                                                       |

## Background

The N-end rule pathway is one proteolytic pathway of the ubiquitin system. The recognition component of this pathway, encoded by this gene, binds to a destabilizing N-terminal residue of a substrate protein and participates in the formation of a substrate-linked multiubiquitin chain. This leads to the eventual degradation of the substrate protein. The protein described in this record has a RING-type zinc finger and a UBR-type zinc finger. Mutations in this gene have been associated with Johanson-Blizzard syndrome. [provided by RefSeq, Jul 2008],developmental stage:Expressed in fetal pancreas.,disease:Defects in UBR1 are a cause of Johanson-Blizzard syndrome (JBS) [MIM:243800]. This disorder includes congenital exocrine pancreatic insufficiency, multiple malformations such as nasal wing aplasia, and frequent mental retardation. Pancreas of individuals with JBS do not express UBR1 and show intrauterine-onset destructive pancreatitis.,domain:The RING-H2 zinc finger is an atypical RING finger with a His ligand in place of the fourth Cys of the classical motif.,function:E3 ubiquitin-protein ligase which is a component of the N-end rule pathway. Recognizes and binds to proteins bearing specific N-terminal residues that are destabilizing according to the N-end rule, leading to their ubiquitination and subsequent degradation. May be involved in pancreatic homeostasis.,pathway:Protein modification; protein ubiquitination.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the UBR1 family.,similarity:Contains 1 RING-type zinc finger.,similarity:Contains 1 UBR-type zinc finger.,subunit:Interacts with RECQL4.,tissue specificity:Broadly expressed, with highest levels in skeletal muscle, kidney and pancreas. Present in acinar cells of the pancreas (at protein level),.


## Research Area

## Image Data




Western blot analysis of lysates from HepG2 and Jurkat cells, using UBR1 Antibody. The lane on the right is blocked with the synthesized peptide.

**Product Name: Ubr1 Rabbit Polyclonal Antibody**  
**Catalog #: APRab19582**



Western blot analysis of the lysates from HUVEC cells using UBR1 antibody.



Western Blot analysis of various cells using Ubr1 Polyclonal Antibody diluted at 1: 2000. Secondary antibody was diluted at 1:20000

## **Note**

For research use only.