

Product Name: VDAC1 Rabbit Polyclonal Antibody**Catalog #: APRab19758**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ELISA 1:10000-1:20000
Molecular Weight	31kDa

Antigen Information

Gene Name	VDAC1 VDAC1; VDAC; Voltage-dependent anion-selective channel protein 1; VDAC-1; hVDAC1;
Alternative Names	Outer mitochondrial membrane protein porin 1; Plasmalemmal porin; Porin 31HL; Porin 31HM
Gene ID	7416.0
SwissProt ID	P21796
Immunogen	The antiserum was produced against synthesized peptide derived from the N-terminal region of human VDAC1. AA range:1-50


Background

This gene encodes a voltage-dependent anion channel protein that is a major component of the outer mitochondrial membrane. The encoded protein facilitates the exchange of metabolites and ions across the outer mitochondrial membrane and may regulate mitochondrial functions. This protein also forms channels in the plasma membrane and may be involved in transmembrane electron transport. Alternate splicing results in multiple transcript variants. Multiple pseudogenes of this gene are found on chromosomes 1, 2 3, 6, 9, 12, X and Y.[provided by RefSeq, Sep 2010],domain:Consists mainly of a membrane-spanning beta-barrel formed by 19 beta-strands. The helical N-terminus folds back into the pore opening and plays a role in voltage-gated channel activity.,function:Forms a channel through the mitochondrial outer membrane and also the plasma membrane. The channel at the outer mitochondrial membrane allows diffusion of small hydrophilic molecules; in the plasma membrane it is involved in cell volume regulation and apoptosis. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV. The open state has a weak anion selectivity whereas the closed state is cation-selective. May participate in the formation of the permeability transition pore complex (PTPC) responsible for the release of mitochondrial products that triggers apoptosis.,similarity:Belongs to the eukaryotic mitochondrial porin family.,subunit:Interacts with hexokinases (By similarity). Interacts with BCL2L1. Interacts with influenza A virus PB1-F2 protein.,tissue specificity:Heart, liver and skeletal muscle.,

Research Area

Calcium;Parkinson's disease;Huntington's disease;

Image Data



Western blot analysis of lysate from K562 cells, using VDAC1 Antibody.