

# **Product Name: VN2R1P Rabbit Polyclonal Antibody**

Catalog #: APRab19816

For research use only.

### **Summary**

**Description** Rabbit polyclonal Antibody

**Host** Rabbit

ApplicationWB,ICC/IF,ELISAReactivityHuman,Rat,MouseConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

ClonalityPolyclonalFormLiquidConcentration1mg/ml

**Storage** Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

**Shipping** Ice bags

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer** 

preservative N.

**Purification** Affinity purification

## **Application**

**Dilution Ratio** WB 1:500-1:2000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000

Molecular Weight 83kDa

## **Antigen Information**

Gene Name CASRL1

**Alternative Names** 

 Gene ID
 344760.0

 SwissProt ID
 Q8NGV9

The antiserum was produced against synthesized peptide derived from human CASRL1. AA Immunogen

range:400-449

## **Background**


The CARL-1 monoclonal antibody reacts with human TWEAK, a type II transmembrane TNF superfamily member with high




identity to TNF in its extracellular portion. TWEAK transcript is expressed broadly in many adult and fetal tissues, however, the staining of human peripheral blood mononuclear cells with monoclonal antibodies shows a more restricted pattern. While freshly isolated PBMCs do not express detectable levels of TWEAK on their surface, IFN-gamma-stimulated blood monocytes rapidly upregulate TWEAK surface expression. TWEAK is expressed as membrane bound and secreted forms. Interaction of TWEAK with its counter-receptor promotes secretion of IL-8, activation of NF-kappaB, proliferation of endothelial cells, and apoptosis in a number of human cell lines. Initially, DR3 was thought to be a receptor for TWEAK, but further studies have shown that TWEAK could induce apoptosis via receptors distinct from DR3. While TWEAK exhibits overlapping signaling functions to TNF, it is generally less effective in inducing apoptosis, giving rise to its name, TNF-like weak inducer of apoptosis. For detection of human TWEAK by sandwich ELISA, a combination of purified CARL-2 for capture and biotinylated CARL-1 for detection is recommended.

#### **Research Area**

# **Image Data**



Immunofluorescence analysis of A549 cells, using CSRL1 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from K562 cells and A549 cells, using CSRL1 Antibody. The lane on the right is blocked with the synthesized peptide.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838