## **Product Name: ZNF225 Rabbit Polyclonal Antibody**

Catalog #: APRab20251



### **Summary**

**Production Name** ZNF225 Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

**Host** Rabbit

**Application** WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA

**Reactivity** Human,Rat,Mouse

#### **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer** 

preservative N.

**Purification** Affinity purification

#### **Immunogen**

Gene Name ZNF225

Alternative Names ZNF225; Zinc finger protein 225

**Gene ID** 7768.0

Q9UK10.The antiserum was produced against synthesized peptide derived from human **SwissProt ID** 

ZNF225. AA range:641-690

## **Application**

**Dilution Ratio** WB 1:500-1:2000, IHC-P 1:100-1:300, ELISA 1:20000, IF-P/IF-F/ICC/IF 1:50-200

Molecular Weight 82kDa

# **Product Name: ZNF225 Rabbit Polyclonal Antibody**

Catalog #: APRab20251




#### **Background**


function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 KRAB domain.,similarity:Contains 18 C2H2-type zinc fingers.,function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 KRAB domain.,similarity:Contains 18 C2H2-type zinc fingers.,

#### Research Area

### **Image Data**



Western blot analysis of lysates from HeLa cells, using ZNF225 Antibody. The lane on the right is blocked with the synthesized peptide.




Western blot analysis of the lysates from HT-29 cells using ZNF225 antibody.


# **Product Name: ZNF225 Rabbit Polyclonal Antibody**

Catalog #: APRab20251





Western Blot analysis of various cells using ZNF225 Polyclonal Antibody. Secondary antibody was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA) .



Immunohistochemical analysis of paraffin-embedded Human thymus gland. Antibody was diluted at 1:100 (4°, overnight) . High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

#### Note

For research use only.