
Product Name: ZNHIT1 Rabbit Polyclonal Antibody**Catalog #: APRab20291**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ELISA 1:10000-1:20000
Molecular Weight	18-22kDa

Antigen Information

Gene Name	ZNHIT1
Alternative Names	ZNHIT1; CGBP1; ZNFN4A1; Zinc finger HIT domain-containing protein 1; Cyclin-G1-binding protein 1; Zinc finger protein subfamily 4A member 1; p18 Hamlet
Gene ID	10467.0
SwissProt ID	O43257
Immunogen	Synthesized peptide derived from ZNHIT1 . at AA range: 40-120

Background

function:Seems to play a role in p53-mediated apoptosis induction.,induction:Induced by DNA damage.,PTM:Phosphorylated

on Thr by MAPK11 or MAPK14.,PTM:Stres-induced ZNHIT1 is mainly regulated at the level of protein.,similarity:Belongs to the ZNHIT1 family.,similarity:Contains 1 HIT-type zinc finger.,subunit:Interacts with MAPK11 and MAPK14. Component of the chromatin-remodeling SRCAP complex composed of at least SRCAP, DMAP1, RUVBL1, RUVBL2, ACTL6A, YEATS4, ACTR6 and ZNHIT1.,function:Seems to play a role in p53-mediated apoptosis induction.,induction:Induced by DNA damage.,PTM:Phosphorylated on Thr by MAPK11 or MAPK14.,PTM:Stres-induced ZNHIT1 is mainly regulated at the level of protein.,similarity:Belongs to the ZNHIT1 family.,similarity:Contains 1 HIT-type zinc finger.,subunit:Interacts with MAPK11 and MAPK14. Component of the chromatin-remodeling SRCAP complex composed of at least SRCAP, DMAP1, RUVBL1, RUVBL2, ACTL6A, YEATS4, ACTR6 and ZNHIT1.,

Research Area

Cell Biology; Apoptosis; Intracellular; p53 Pathway

Image Data



Western Blot analysis of various cells using ZNHIT1 Polyclonal Antibody diluted at 1 : 1000. Secondary antibody was diluted at 1:20000