Product Name: **BMP4 Rabbit Monoclonal antibody**
Catalog #: **AMRe01731**
For research use only.

Summary

Description	Recombinant rabbit monoclonal antibody
Host	Rabbit
Application	WB,ICC/IF,IP
Reactivity	Human
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal Antibody
Form	Liquid
Concentration	0.15mg/ml. The concentration of this product may be batch-dependent.
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% protective protein
Purification	Affinity Purified

Application

Dilution Ratio	WB 1:500-1:1000,ICC/IF 1:50-1:200,IP 1:20-1:50
Molecular Weight	Calculated MW: 47 kDa; Observed MW: 47 kDa

Antigen Information

Gene Name	BMP4
Alternative Names	BMP4; BMP2B; DVR4; Bone morphogenetic protein 4; BMP-4; Bone morphogenetic protein 2B; BMP-2B
Gene ID	652
SwissProt ID	P12644
Immunogen	A synthetic peptide of human BMP4

Background

Bone morphogenetic proteins (BMPs) were first identified as molecules that can induce ectopic bone and cartilage formation.

BMPs belongs to the TGF-β superfamily, playing many diverse functions during development. BMPs are synthesized as precursor proteins and then processed by cleavage to release the c-terminal mature BMP. BMPs initiate signaling by binding to a receptor complex containing type I and type II serine/threonine receptor kinases that then phosphorylate Smad (mainly Smad1, 5 and 8), resulting the translocation of Smad into the nucleus. BMP was also reported to activate MAPK pathways in some systems.

Research Area

Cardiovascular

Image Data



Immunocytochemistry analysis of BMP4 (green) in HEPG2 using BMP4 antibody,and DAPI(blue).



Western blot analysis of BMP4 in K562, Hela lysates using BMP4 antibody.