# **Product Name: CPSF6 Rabbit Monoclonal Antibody**

Catalog #: AMRe01849



### **Summary**

Production Name CPSF6 Rabbit Monoclonal Antibody

**Description** Rabbit Monoclonal antibody

**Host** Rabbit

**Application** WB,IHC-F,IHC-P,ICC/IF,IP

**Reactivity** Human, Mouse, Rat

#### **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

Clonality Monoclonal Form Liquid

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05%

protective protein

**Purification** Affinity Purification

#### **Immunogen**

Buffer

Gene Name CPSF6

Alternative Names CFIM; CFIM68; CFIM72; HPBRII-4; HPBRII-7

 Gene ID
 11052

 SwissProt ID
 Q16630.

# **Application**

**Dilution Ratio** WB: 1:500-1:1000 IHC: 1:50-1:100 IF: 1:50-1:200 IP: 1:20

Molecular Weight Calculated MW: 59 kDa; Observed MW: 70 kDa

## **Background**

Component of the cleavage factor Im (CFIm) complex that functions as an activator of the pre-mRNA 3'-end cleavage and

# Product Name: CPSF6 Rabbit Monoclonal Antibody Catalog #: AMRe01849




polyadenylation processing required for the maturation of pre-mRNA into functional mRNAs (PubMed:9659921, PubMed:8626397, PubMed:14690600, PubMed:29276085). CFIm contributes to the recruitment of multiprotein complexes on specific sequences on the pre-mRNA 3'-end, so called cleavage and polyadenylation signals (pA signals) (PubMed:9659921, PubMed:8626397, PubMed:14690600). Most pre-mRNAs contain multiple pA signals, resulting in alternative cleavage and polyadenylation (APA) producing mRNAs with variable 3'-end formation (PubMed:23187700, PubMed:29276085). The CFIm complex acts as a key regulator of cleavage and polyadenylation site choice during APA through its binding to 5'-UGUA-3' elements localized in the 3'-untranslated region (UTR) for a huge number of pre-mRNAs (PubMed:20695905, PubMed:29276085). CPSF6 enhances NUDT21/CPSF5 binding to 5'-UGUA-3' elements localized upstream of pA signals and promotes RNA looping, and hence activates directly the mRNA 3'-processing machinery (PubMed:15169763, PubMed:29276085, PubMed:21295486). Plays a role in mRNA export (PubMed:19864460).

#### **Research Area**

**Epigenetics and Nuclear Signaling** 

### **Image Data**



Western blot analysis of CPSF6 in rat Brain, MCF-7, 3T3, C6 lysates using CPSF6 antibody.

#### Note

For research use only.