

Product Name: Hydroxyl-Histone H2A (Tyr39) Rabbit Monoclonal Antibody**Catalog #: AMRe03278**

For research use only.

Summary

| | |
|----------------------|--|
| Description | Recombinant rabbit monoclonal antibody |
| Host | Rabbit |
| Application | WB,IHC |
| Reactivity | Human,Mouse |
| Conjugation | Unconjugated |
| Modification | Hydroxylated |
| Isotype | IgG |
| Clonality | Monoclonal |
| Form | Liquid |
| Concentration | 0.5mg/ml. The concentration of this product may be batch-dependent. |
| Storage | Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles. |
| Shipping | Ice bags |
| Buffer | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% protective protein |
| Purification | Affinity Purification |

Application

| | |
|-------------------------|--|
| Dilution Ratio | WB 1:500-1:1000,IHC 1:50-1:100 |
| Molecular Weight | Calculated MW: 14 kDa; Observed MW: 14 kDa |

Antigen Information

| | |
|--------------------------|---|
| Gene Name | H2AC4 H2A.1; H2A/c; H2A1; H2AFC.; H2AFD; H2AFI; H2AFN; H2AFP; HIST1H2AG; HIST1H2AI; |
| Alternative Names | HIST1H2AK; HIST1H2AL; HIST1H2AM; histone cluster 1; H2ai; Histone H2A type 1; Histone H2A/p |
| Gene ID | 3012 |
| SwissProt ID | P04908 |
| Immunogen | A synthetic Hydroxylated peptide corresponding to residues target protein |


Background

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.


Research Area

Epigenetics and Nuclear Signaling


Image Data



Western blot analysis of Hydroxyl-Histone H2A (Tyr39) in 3T3 lysates using Hydroxyl-Histone H2A (Tyr39) antibody.



Western blot analysis of Hydroxyl-Histone H2A (Tyr39) in HeLa, A549, HL-60 lysates using Hydroxyl-Histone H2A (Tyr39) antibody. Observed band size: 14 kDa.



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using Hydroxyl-Histone H2A (Tyr39) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.